



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 0 998 850 A1**

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
10.05.2000 Bulletin 2000/19

(51) Int. Cl.⁷: **A01N 31/02**

(21) Application number: **99121907.2**

(22) Date of filing: **05.11.1999**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **06.11.1998 JP 31573798**

(71) Applicant: **Kao Corporation**
Chuo-Ku Tokyo 103 (JP)

(72) Inventors:
• **Hayashi, Masaharu**
Wakayama-shi, Wakayama (JP)

• **Suzuki, Tadayuki**
Wakayama-shi, Wakayama (JP)
• **Kamei, Masatoshi**
Wakayama-shi, Wakayama (JP)
• **Hayashi, Toshio**
Wakayama-shi, Wakayama (JP)
• **Kurita, Kazuhiko**
Wakayama-shi, Wakayama (JP)

(74) Representative: **HOFFMANN - EITLE**
Patent- und Rechtsanwälte
Arabellastrasse 4
81925 München (DE)

(54) **Plant-activating agent**

(57) The present invention provides a plant-activating agent which effects no chemical injury on plant and which makes an activity of plant improve efficiently. That is to say, the present invention provides a method of activating a plant, which comprises treating the plant with a mono-alcohol having 12 to 24 carbon atoms as a plant-activating agent; use of the mono-alcohol as a plant-activating agent; and a plant-activating agent or composition comprising the mono-alcohol; optionally with a surfactant, a fertilizer component and/or a chelating agent.

EP 0 998 850 A1

Description

TECHNICAL FIELD

5 [0001] The present invention relates to a plant-activating agent and/or a plant-activating composition. Also, as use thereof, the present invention provides a method of activating a plant by applying thereof in the state of solution or solid to a root, stem, phylloplane or fruit of plant, such as spraying onto phylloplanes and irrigating into soil. Now, hereinafter, the term of "plant" includes plants, vegetables, fruits, crops, seeds, flowers, herbs, floras, and so on.

10 BACKGROUND ART

[0002] Various nutrient elements are necessary for growth of plants. It is known that lack of some of the elements causes the hindrance of the growth of the plants. For example, the big three fertilizer components function as follows. Nitrogen is a component element of proteins, and phosphorus is a formation element of nucleic acid or phosphorus lipid and further plays an important part in energy metabolism and synthetic or decomposing reaction of a substance. Potas-
 15 sium has a physiological action of substance metabolism or substance migration. If these main components lacks, the growth of plants generally becomes poor. Calcium is an important component constituting the plant-realities and cells, and further plays an important part in maintenance of the balance of the metabolic system. The lacking state of calcium causes physiological troubles. Besides, various nutrients as follows are necessary for plants: Mg, Fe, S, B, Mn, Cu, Zn, Mo, Cl, Si, Na and the like.

[0003] Nutritious components such as nitrogen, phosphorus and potassium are applied as basal fertilizer or additional fertilizer. Alternatively, they are applied by diluting liquid fertilizer and irrigating the diluted fertilizer into soil or by spraying the diluted fertilizer onto phylloplanes. These fertilizers are necessary and/or essential for the growth of plants. However, even if they are applied at larger concentrations than some values, the growth of plants and the yield of the
 25 plants cannot be further improved.

[0004] However, it is an important theme in agricultural production to promote the growth of agricultural plants and increase the yield per unit area to strive for an increase in income. Various plant growth regulators being necessary for this have been developed and used. The plant growth regulators, the typical examples of which include gibberellin and auxin, are used to regulate growth reaction or form-producing reaction such as germination, rooting, expansion, flower-
 30 ing and bearing. The actions of these substances are many-sided or complicated. The uses thereof are restrictive.

[0005] In order to solve such problems, there are known a phylloplane spraying agent using an oligosaccharide (JP-A 9-322647), and techniques in which a liquid fertilizer comprising a sugar, a mineral, an amino acid, an extract from seaweeds, or a fermentation extract of microorganisms is sprayed onto phylloplanes or is applied in the form of the liquid. In the present situation, however, their effects are insufficient for practical use.

[0006] For an increased yield, when a large amount of fertilizer is applied into the soil, various components may become excessive in the soil so that the balance of absorption thereof may become bad or the growth of plants may be delayed. As a result, there arise, for example, problems that the increased yield, as an aim, cannot be attained or the quality such as sugar concentration (Brix. value) or freshness (green degree) does not rise. And then, since there are a limit of absorption from roots which aim absorption of nutrients, direct absorption of necessary fertilizer elements from
 40 phylloplanes or fruits is attempted by spraying an aqueous solution or aqueous suspension of the elements. However, even if the aqueous solution of the necessary elements is merely sprayed onto phylloplanes, a problem arises from the viewpoint of absorption efficiency. Spraying excessive amounts of fertilizer elements impose stress on plants, resulting chemical injury.

[0007] Handbook of Agricultural Chemicals (edited in 1994), on page 475, discloses decyl alcohol as a restrainer of an axillary bud of a tobacco-plant. JP-A 55-40674 discloses an alcohol having 30 carbon atoms as a plant growth promoter.

[0008] It is known to dilute a foam concentrated product with water, generate foams at a static pressure of 15 psi or more in a foam generator connected to a tap water pipe, and treat plants or soil with the resultant foams (US-A 3922927). However, this patent never discloses nor suggests use as a plant-activating agent or a method for activating
 50 plants.

DISCLOSURE OF THE INVENTION

[0009] The present invention relates to a plant-activating agent comprising a mono-alcohol having 12-24 carbon atoms and relates to a plant-activating composition comprising the mono-alcohol, and a surfactant, a fertilizer component or a chelating agent.

[0010] Namely, the present invention provides a plant-activating agent comprising a mono-alcohol having 12-24 carbon atoms, and provides a plant-activating composition comprising the mono-alcohol and at least one compound

selected from a surfactant and a chelating agent.

[0011] The invention provides a method of activating a plant, which comprises treating the plant with a mono-alcohol having 12 to 24 carbon atoms as a plant-activating agent; and use of the mono-alcohol as a plant-activating agent.

[0012] It is preferred that the plant is treated further with at least one compound selected from a surfactant, a fertilizer component and a chelating agent, in use or method of the above-mentioned.

[0013] The surfactant may be selected from an ester group-containing nonionic surfactant, an ether group-containing nonionic surfactant having no nitrogen atom, an amphoteric surfactant, a carboxylic anionic surfactant and a phosphoric anionic surfactant, in use or method of the above-mentioned.

[0014] The invention provides a plant-activating composition comprising a mono-alcohol having 12 to 24 carbon atoms and one group selected from (i) a chelating agent, (ii) a surfactant and a chelating agent, (iii) a fertilizer component and a chelating agent, and (iv) a surfactant, a fertilizer component and a chelating agent.

[0015] Another composition may preferably comprise a mono-alcohol having 12 to 24 carbon atoms and at least one surfactant selected from an ester group-containing nonionic surfactant, a ether group-containing nonionic surfactant having no nitrogen atom, an amphoteric surfactant, a carboxylic anionic surfactant and a phosphoric anionic surfactant.

[0016] A plant-activating composition of the invention comprises a mono-alcohol having 12 to 24 carbon atoms and at least one surfactant selected from an ester group-containing nonionic surfactant, an ether group-containing nonionic surfactant having no nitrogen atom, an amphoteric surfactant, a carboxylic anionic surfactant and a phosphoric anionic surfactant, optionally further comprising a fertilizer component.

[0017] A plant-activating composition of the invention comprises a mono-alcohol having 12 to 24 carbon atoms and one group selected from (iii) a fertilizer component and a chelating agent, (iv) a surfactant, a fertilizer component and a chelating agent and (v) a surfactant and a fertilizer component.

MODES FOR CARRYING OUT THE INVENTION

[0018] In the present invention, a mono-alcohol having 12-24 carbon atoms, preferably 14-22 carbon atoms, particularly preferably 16-20 carbon atoms, is used since the alcohol can effectively give plant vital power activation without causing chemical injury. The hydrocarbon group of the mono-alcohol may be saturated or unsaturated, and may be in the form of a straight chain, a branched chain or a cyclic chain. The hydrocarbon group is preferably an alkyl group being a straight or branched chain, particularly preferably an alkyl group being a straight chain. Specific examples of the mono-alcohol include lauryl alcohol, cetyl alcohol, stearyl alcohol, eicosanol, behenyl alcohol, phytol, oleyl alcohol, and alcohols originated from natural fats and/or oils.

[0019] The form of the plant-activating agent comprising the mono-alcohol according to the present invention may be any form, such as a liquid, a flowable, a paste, a wettable powder, a granule, a dust formulation or a tablet. At the time of use, the plant-activating agent is generally sprayed in the form of an aqueous solution, an aqueous dispersion, or an aqueous emulsion which has a mono-alcohol concentration of from 1 to 500 ppm onto phylloplanes or roots of a plant.

[0020] In order to supply the plant-activating agent of the present invention to a plant, various methods may be used. Examples of the methods include a method of applying a dust formation or a granule as fertilizer directly, a method of spraying a diluted aqueous solution directly onto phylloplanes, stems or fruits of a plant, a method of injecting a diluted aqueous solution into soil, and a method of supplying to dilute and to mix into a liquid for a hydroponics and a supplying water which are contacted with roots and which are such as a hydroponics and a rock wool.

[0021] Plants, which can be treated with the plant-activating agent of the present invention, may be a fruit vegetable such as a cucumber, a pumpkin, a watermelon-plant, a melon, a tomato, an eggplant, a green pepper, a strawberry, an okra, kidney beans in a pod, a broad bean, a pea, green soybeans in a pod and a corn; a leaf vegetables such as a Chinese cabbage, greens for pickling, a Brassica campestris (a Chinese spinach-like green vegetable), a cabbage, a cauliflower, a broccoli, a Brussels sprout, an onion, a Welsh onion, a garlic, a scallion, a leek, an asparagus, a lettuce, a green for salad (which is called Saladana in Japan), a celery, a spinach, a crown daisy, a parsley, a trefoil (which is called Mitsuba in Japan and is useful as herb), a dropwort, an udo (which is an Aralia cordata), a Japanese ginger, a Japanese butterbur and a labiate; and a root vegetable such as a radish, a turnip, a burdock, a carrot, a potato, a taro, a sweet potato, a yam, a ginger-plant (which is called Shoga in Japan) and a lotus root. Besides, the plant-activating agent may be used for a rice-plant; a barley, a wheat or a group thereof; petalous-plants and the like.

[0022] In the present invention, the following surfactant is preferably used together with the above-mentioned mono-alcohol to promote emulsification, dispersion, solubilization and permeation of the mono-alcohol.

[0023] Examples of nonionic surfactants include sorbitan fatty acid esters, polyoxyalkylene sorbitan fatty acid esters, polyoxyalkylene fatty acid esters, glycerol fatty acid esters, polyoxyalkylene glycerol fatty acid esters, polyglycerol fatty acid esters, polyoxyalkylene polyglycerol fatty acid esters, sucrose fatty acid esters, resin acid esters, polyoxyalkylene resin acid esters, polyoxyalkylene alkyl ethers, polyoxyalkylene alkylphenyl ethers, alkyl(poly)glycosides and

polyoxyalkylenealkyl(poly)glycosides. Preferably, an ether group-containing nonionic surfactant having no nitrogen atom and ester group-containing nonionic surfactant may be cited.

[0024] Examples of anionic surfactants include carboxylic, sulfonic, sulfuric ester group-containing and phosphoric ester group-containing surfactants, and a carboxylic and phosphoric ester group-containing surfactants are preferred.

[0025] Examples of the carboxylic surfactants include fatty acids having 6-30 carbon atoms or salts thereof, polyhydric carboxylic acid salts, polyoxyalkylene alkyl ether carboxylic acid salts, polyoxyalkylene alkylamide ether carboxylic acid salts, rhodinic acid salts, dimmer acid salts, polymer acid salts and tall oil fatty acid salts.

[0026] Examples of the sulfonic surfactants include alkylbenzenesulfonic acid salts, alkylsulfonic acid salts, alkyl-naphthalenesulfonic acid salts, naphthalenesulfonic acid salts, diphenyl ether sulfonic acid salts, condensates of alkyl-naphthalenesulfonic acid, and condensate of naphthalenesulfonic acid.

[0027] Examples of the sulfuric ester group-containing surfactants include alkylsulfuric ester salts (alkylsulfuric acid salts), polyoxyalkylene alkylsulfuric ester salts (polyoxyalkylene alkylsulfuric acid salts), polyoxyalkylene alkyl phenyl ether sulfuric acid salts, tristyrenated phenol sulfuric acid ester salts, polyoxyalkylene distyrenated phenol sulfuric acid ester salts and alkylpolyglycoside sulfuric acid salts.

[0028] Examples of phosphoric acid ester group-containing surfactants include alkyl phosphoric acid ester salts, alkylphenylphosphoric acid ester salts, polyoxyalkylene alkylphosphoric acid ester salts and polyoxyalkylene alkyl-phenylphosphoric acid ester salts.

[0029] Examples of the salts include metallic salts (such as salts of Na, K, Ca, Mg and Zn), ammonium salts, alkanol amine salts and aliphatic amine salts.

[0030] Examples of amphoteric surfactants include amino acid group-containing, betaine group-containing, imidazoline group-containing and amine oxide group-containing surfactants.

[0031] Examples of the amino acid group-containing surfactants include acylamino acid salts, acylsarcosine acid salts, acyloylmethylaminopropionic acid salts, alkylaminopropionic acid salts and acylamide ethylhydroxyethylmethyl-carboxylic acid salts.

[0032] Examples of the betaine group-containing surfactants include alkyltrimethylbetaine, alkylhydroxyethylbetaine, acylamide propylhydroxypropylammonia sulfobetaine, acylamide propylhydroxypropyl ammonia sulfobetaine and ricinoleic acid amide propyl dimethylcarboxy methylammonia betaine.

[0033] Examples of the imidazoline group-containing surfactants include alkylcarboxy methylhydroxy ethylimidazolinium betaine and alkylethoxy carboxy methylimidazolinium betaine.

[0034] Examples of the amine oxide group-containing surfactants include alkyltrimethylamine oxide, alkyl-diethanolamine oxide and alkylamidepropylamine oxide.

[0035] One kind of the above-mentioned surfactants may be used, and a mixture of two or more kinds of the above-mentioned surfactants may be used. In the case that one of these surfactants comprises a polyoxyalkylene group, the polyoxyalkylene group is preferably a polyoxyethylene group and the average mole number of added polyoxyethylene groups is preferably from 1 to 50.

[0036] As the surfactant, at least one compound selected from ester group-containing nonionic surfactants, ether group-containing nonionic surfactants having no nitrogen atom, amphoteric surfactants, carboxylic anionic surfactants and phosphoric anionic surfactant is preferable.

[0037] In particular, it is preferable to be at least one compound selected from ester group-containing nonionic surfactants and ether group-containing nonionic surfactants having no nitrogen atom.

[0038] The following fertilizer components may be used together with the above-mentioned mono-alcohol. Specific examples thereof may be inorganic or organic compounds which can supply elements such as N, F, K, Ca, Mg, S, B, Fe, Mn, Cu, Zn, Mo, Cl, Si and Na, in particular N, P, K, Ca and Mg. Examples of such inorganic compounds include ammonium nitrate, potassium nitrate, ammonium sulfate, ammonium chloride, ammonium phosphate, sodium nitrate, urea, ammonium carbonate, potassium phosphate, calcium superphosphate, fused phosphate fertilizer ($3\text{MgO} \cdot \text{CaO} \cdot \text{P}_2\text{O}_5 \cdot 3\text{CaSiO}_2$), potassium sulfate, potassium chloride, nitrate of lime, slaked lime, carbonate of lime, magnesium sulfate, magnesium hydroxide and magnesium carbonate. Examples of the organic compounds include fowl droppings, cow dung, Bark compost, amino acid, peptone, amino acid solution (which is called Miei in Japan), fermentation extracts, calcium salts of organic acids (such as citric acid, gluconic acid and succinic acid), and calcium salts of fatty acids (such as formic acid, acetic acid, propionic acid, caprylic acid, capric acid and caproic acid). These fertilizer components may be used together with the surfactant. In the case that fertilizer components are sufficiently applied as basal fertilizer to soil as seen in outdoor cultivation of a rice-plant or vegetables, it is unnecessary to mix the fertilizer components. Further, when a cultivation form is such as a fertigation (a hydroponic soil culture) or a hydroponics, when it avoids applying excessively basal fertilizer and when it is a type of providing a fertilizer component as well as irrigation-water, the fertilizer component is preferably mixed.

[0039] When the plant-activating composition of the present invention is mixed with the following organic acid having chelating ability or a salt thereof, the growth and absorption efficiency of fertilizer improved further. Specific examples thereof include oxycarboxylic acids such as citric acid, gluconic acid, malic acid, heptonic acid, oxalic acid, malonic

acid, lactic acid, tartaric acid, succinic acid, fumaric acid, maleic acid, adipic acid and glutaric acid; polyhydric carboxylic acids; and salts thereof such as potassium salt, sodium salt, alkanolamine salt and aliphatic amine salt.

[0040] Mixing a chelating agent besides the organic acids also causes the growth and absorption efficiency of fertilizer to be improved. Examples of the mixed chelating agents include aminocarboxylic group-containing chelating agents such as EDTA, NTA and CDTA.

[0041] The form, spraying method and the like of plant-activating composition of the present invention are the same as described above. If necessary, water and/or a solvent may be added to the composition.

[0042] In the plant-activating composition or the fertilizer composition of the present invention, it is preferable to use respectively; per 100 parts by weight of the mono-alcohol; 10-20000 parts, in particular 100-2000 parts, by weight of the surfactant; 0-50000 parts, in particular 10-5000 parts, by weight of the fertilizer component; 0-1000 parts, in particular 10-500 parts, by weight of the chelating agent; and 0-5000 parts, in particular 10-500 parts, by weight of other nutritious sources (such as sugars, amino acids and vitamins).

[0043] In the case that the plant-activating composition is applied in the form of a dust formulation or granule as fertilizer to soil, it is preferable that the used dust formulation or granule comprise the above-mentioned components except water at the same ratios as above, in general. This dust formulation or granule may comprise a vehicle to prevent caking.

[0044] The plant-activating agent of the present invention makes it possible to improve the activity of plant effectively without causing any chemical injury on plants if the plants are treated with an appropriate concentration of the plant-activating agent. For this reason, it can be used for various plants. According to the present invention, promotion of taking root of a plant, increase of chlorophyll value (SPAD value), increase of absorption efficiency of fertilizer and so on are seen as improvement at growth of plant.

Examples

Example 1 (Test of soil-treatment for tomato seedlings)

[0045]

Species: "Home Momotaro" (Takii & Company LTD)

Vessel for cultivation: a cell tray having 50 holes for germination, and a pot (diameter: 14 cm) for cultivation

Used soil: Takii soil for seeding [N: P₂O₅: K₂O= 480: 750: 345 (mg/L), pH 6.4, and EC: 0.96]

[0046] Under the above-mentioned conditions, seeds were sown in the cell tray having 50 holes. When 2 weeks went by after germination, the plants were transplanted to the pot. At intervals of 1 week after 3 days from the transplantation, the soil was treated 4 times with fertilizer compositions being blended raw materials shown in Table 1 and 1000 ppm of a fertilizer component "Otsuka OKF2" (made by Otsuka Chemical Co., Ltd.). The concentrations of the blended raw materials are shown in Table 1 and their balance was water. The amount for each of the treatments was about 50 ml per pot. The fertilizer compositions were caused to penetrate into the soils. For the respective compositions, the test was repeated 3 times. On the 7th day after the fourth treatment, the growth state [height and weight (,hereinafter, "weight" means raw-weight,) of the respective plants and the SPAD value (SPAD 502 made by Minolta Co., Ltd.) representing a green degree were measured. The measured values obtained from the 3 repetitions were averaged. The averaged value was relatively compared as the value that was obtained from the no-treated area (Comparative product 1-4) and made to be 100. The results are shown in Table 1.

Table 1

	No.	Blended raw materials	Concentration (ppm)	Results of the test		
				Height	Weight	SPAD value
Invention product	1 - 1	Stearyl alcohol	100	148	142	122
		POE(20) sorbitan monooleate	500			
	1 - 2	Eicosanol	200	145	140	119
		POE(20) hardened castor oil	1000			
	1 - 3	Decyldodecanol	100	134	124	110
		POE(15) glycerol beef tallow fatty acid ester	500			
	1 - 4	Lauryl alcohol	50	132	122	112
		Na salt of POE(3) lauryl sulfuric ester	100			
		POE(20) sorbitan monolaurate	200			
	1 - 5	Oleyl alcohol	300	138	130	118
		POE(20) sorbitan monolaurate	600			
		Sorbitan monolaurate	400			
	1 - 6	Stearyl alcohol	200	144	138	122
		Cetyl alcohol	100			
		POE(20) sorbitan monolaurate	500			
		Tetraglycerol monooleate	500			
	1 - 7	Branched C ₁₂ /C ₁₃ alcohol *	100	118	110	112
		POE(10) oleic acid ester	500			
		Alkylglucoside **	200			
	1 - 8	C ₁₆ /C ₁₈ alcohol (weight ratio = 3/7. This is true hereinafter.)	200	158	144	120
		POE(20) sorbitan monooleate	200			
		Sucrose stearic acid ester	300			
	1 - 9	Stearyl alcohol	50	156	147	129
		POE(20) sorbitan monooleate	200			
		EDTA·4Na	20			
	1 - 10	Coconut alcohol (C ₁₄ /C ₁₆ /C ₁₈ = 10/70/20)	100	122	124	118
		Oleic acid monodiglyceride	200			
		POE(20) sorbitan monooleate	500			
Comparative product	1 - 1	Ethanol	200	95	98	99
		POE(20) sorbitan monooleate	200			
	1 - 2	Isopropanol	100	86	98	88
		Na salt of POE(3) lauryl sulfuric ester	100			
		POE(20) sorbitan monolaurate	200			
	1 - 3	Octyl alcohol	100	75	88	86
		POE(20) sorbitan monolaurate	200			
	1 - 4	Water (no-treated area)	—	100	100	100

*: Synthetic C₁₂/C₁₃ alcohol (branch ratio = 30%. This is true hereinafter.)

**: Alkyl(C₁₀/C₁₂/C₁₄ = 60/30/10) polyglucoside

(Notes) In Table 1, polyoxyethylene is abbreviated to POE, the respective numbers inside parentheses are the average mole numbers of added ethylene oxide (the hereinafter is the same as this). Also, for example, notes of C₁₆ and C₁₈ mean the number of carbon atoms being 16 and 18, respectively (the hereinafter is the same as this).

[0047] From the results shown in Table 1, it is noted that the plant-activating composition of the present invention made the growth of the plants promote remarkably and the green degree raise obviously, compared with comparative products.

Example 2 (Test of soil-treatment for Brassica campestris seedlings)

[0048]

- 5 Brassica campestris seeds: Takii & Company LTD
Vessel for cultivation: a cell tray having 50 holes
Used soil: Takii soil for seeding (,which was the same as in Example 1)

10 [0049] Under the above-mentioned conditions, seeds were sown in the cell tray having 50 holes. At intervals of 1
week after 2 weeks from germination, the soil was treated 4 times with fertilizer compositions being blended raw mate-
rials shown in Table 2 and 1000 ppm of a fertilizer component "Otsuka OKF2". The concentrations of the blended raw
materials are shown in Table 2 and their balance was water. The amount for each of the treatments was about 60 ml for
respective 10 holes. The fertilizer compositions were caused to penetrate into the soils. For the respective composi-
tions, the test for 10 holes was performed 3 times, i.e. 3 repetitions. On the 7th day after the fourth treatment, the growth
15 state (height and weight) of the respective plants and the SPAD value (SPAD 502 made by Minolta Co., Ltd.) represent-
ing a green degree were measured. The measured values obtained from the 3 repetitions were averaged. The aver-
aged value was relatively compared as the value that was obtained from the no-treated area (Comparative product 2-
4) and was made to be 100. The results are shown in Table 2.

20

25

30

35

40

45

50

55

Table 2

	No.	Blended raw materials	Concentration (ppm)	Results of the test		
				Height	Weight	SPAD value
Invention product	2-1	Lauryl alcohol	100	142	138	110
		POE(20) sorbitan monooleate	200			
	2-2	Myristyl alcohol	100	148	140	114
		POE(20) sorbitan monooleate	200			
	2-3	Cetyl alcohol	100	158	144	114
		POE(20) sorbitan monooleate	200			
	2-4	Stearyl alcohol	100	160	152	140
		POE(20) sorbitan monooleate	200			
	2-5	Oleyl alcohol	100	152	160	138
		POE(20) sorbitan monooleate	200			
	2-6	Eicosanol	100	140	138	120
		POE(20) sorbitan monooleate	200			
	2-7	Behenyl alcohol	100	132	142	118
		POE(20) sorbitan monooleate	200			
	2-8	Phytol	100	142	146	110
		POE(20) sorbitan monooleate	200			
	2-9	C ₁₆ /C ₁₈ alcohol(C ₁₆ /C ₁₈ =3/7)	100	156	162	136
		POE(20) sorbitan monooleate	200			
	2-10	C ₁₆ /C ₁₈ alcohol(C ₁₆ /C ₁₈ =3/7)	100	168	176	136
		POE(20) sorbitan monooleate	200			
	2-11	Heptononic acid Na salt	20	166	176	132
		C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ =3/7)	100			
	2-12	POE(20) sorbitan monooleate	200	172	172	130
		Malonic acid	20			
	2-13	C ₁₆ /C ₁₈ alcohol(C ₁₆ /C ₁₈ =3/7)	100	136	148	112
		POE(20) sorbitan monooleate	200			
	2-14	Maleic acid	20	146	156	118
		Branched C ₁₂ /C ₁₃ alcohol	100			
Comparative product	2-1	POE(20) sorbitan monooleate	200	96	86	90
		Ethanol	100			
	2-2	POE(20) sorbitan monooleate	200	74	68	88
		Octyl alcohol	100			
	2-3	POE(20) sorbitan monooleate	200	82	76	88
		Decyl alcohol	100			
	2-4	POE(20) sorbitan monooleate	200	100	100	100
		Water (no-treated area)	—			

Example 3 (Test of phylloplane-treatment for rice-plant seedlings)

[0050]

- 5 Species: Kinuhikari
 Vessel for cultivation: a cell tray having 50 holes
 Used soil: Ryujou Iseki Baido (seedling soil for rice-plant) (N:P:K = 1:1:1) (made by Iseki & Company, LTD)

10 [0051] Under the above-mentioned conditions, seeds were sown in the cell tray having 50 holes. After the plants
 germinated, the plants fell in the dark age followed by the virescent age. At a period when the height of the plants
 became about 3 cm (after 3 days), the phylloplanes were treated 1 time with fertilizer compositions being blended raw
 materials shown in Table 3 and 1000 ppm of a fertilizer component "Otsuka OKF2". The concentrations of the blended
 raw materials are shown in Table 3 and their balance was water. The amount for each of the treatments was about 1 L
 15 per 50 holes. At the 3.2 leaf age, which was final during the raising seedling period, the growth state (height, weight,
 weight of the portion on the ground and weight of the portion under the ground) of the respective plants and the SPAD
 value (SPAD 502 made by Minolta Co., Ltd.) representing a green degree of a leaf were measured. The measured value
 was relatively compared as the value that was obtained from the no-treated area (Comparative product 3-4) and made
 to be 100. The results are shown in Table 3.

20

25

30

35

40

45

50

55

Table 3

	No.	Blended raw materials	Concentration (ppm)	Results of the test			
				Height n=20	weight of the portion on the ground n=20	weight of the portion under the ground n=20	SPAD value n=20
Invention product	3-1	Cetyl alcohol	10	140	106	112	110
		POE(20) sorbitan monooleate	50				
	3-2	Cetyl alcohol	50	144	108	116	114
		POE(20) sorbitan monooleate	250				
	3-3	Cetyl alcohol	100	152	114	120	120
		POE(20) sorbitan monooleate	500				
	3-4	Cetyl alcohol	500	156	116	130	128
		POE(20) sorbitan monooleate	1000				
	3-5	Stearyl alcohol	10	138	108	120	116
		POE(20) sorbitan monooleate	50				
	3-6	Stearyl alcohol	50	142	110	124	120
		POE(20) sorbitan monooleate	250				
Comparative product	3-1	Octyl alcohol	10	90	92	96	86
		POE(20) sorbitan monooleate	50				
	3-2	Octyl alcohol	50	88	80	74	76
		POE(20) sorbitan monooleate	250				
	3-3	Octyl alcohol	100	68	72	62	66
		POE(20) sorbitan monooleate	500				
	3-4	Water (no-treated area)	—	100	100	100	100
	3-1	Octyl alcohol	10	90	92	96	86
		POE(20) sorbitan monooleate	50				
	3-2	Octyl alcohol	50	88	80	74	76
		POE(20) sorbitan monooleate	250				
	3-3	Octyl alcohol	100	68	72	62	66
		POE(20) sorbitan monooleate	500				
	3-4	Water (no-treated area)	—	100	100	100	100

Example 4 (Test of reproductive ability using chlorella cells)

[0052] Chlorella cells, which are green cells of a higher plant, were cultured with vibration in an inorganic-salt medium. Raw materials shown in Table 4 were added thereto, and then reproductive ability of chlorella cells (ability for increasing the number of the cells) was evaluated by comparison with the result obtained from a medium which was no-treated (i.e., only original nutrients of the inorganic salt medium). For the respective compositions, the test was repeated 3 times. The concentration of the cells was set to 1.00×10^5 cells (per ml) at the start of the test. The numbers of the chlorella cells after 10 days from the addition of the various raw materials followed by culture of the cells are shown as the relative values compared with the value that was obtained from the medium which was no-treated (Comparative product 4-9) and made to be 100. The results are shown in Table 4. As the inorganic salt medium, a Linsmaier-Skoog (LS) medium was used.

Table 4

	Test No.	Blended raw materials	Concentration (ppm)	Result of the test
				proliferating ability of cells
Invention product	4-1	Lauryl alcohol	10	115
	4-2	Myristyl alcohol	10	128
	4-3	Myristyl alcohol	5	138
		Stearyl alcohol	5	
	4-4	Cetyl alcohol	10	148
	4-5	Cetyl alcohol	5	160
		Stearyl alcohol	5	
	4-6	Stearyl alcohol	10	168
	4-7	Oleyl alcohol	10	154
	4-8	Eicosanol	10	152
	4-9	Docosanol	10	130
	4-10	Tetracosanol	10	122
	4-11	Lauryl alcohol	10	119
		Malonic acid	4	
	4-12	Cetyl alcohol	10	152
		Ascorbic acid Na salt	4	
	4-13	Stearyl alcohol	10	172
		EDTA·4Na	4	
	4-14	Eicosanol	10	160
		EDTA·4Na	4	
	4-15	Cetyl alcohol	30	142
Comparative product	4-16	Stearyl alcohol	30	156
	4-17	Eicosanol	30	146
	4-18	Cetyl alcohol	30	146
		Ascorbic acid Na salt	15	
	4-19	Stearyl alcohol	30	159
		EDTA·4Na	15	
	4-20	Eicosanol	30	150
		EDTA·4Na	15	
	4-21	Tetracosanol	10	128
		Fumaric acid	4	
	4-1	Propanol	10	96
	4-2	Hexanol	10	93
	4-3	Propanol	5	92
		Octanol	5	
	4-4	Decyl alcohol	10	91
	4-5	Decyl alcohol	10	94
		Fumaric acid	4	
	4-6	Hexanol	5	96
		Ethanol	5	
	4-7	Octanol	30	95
	4-8	Propanol	10	98
		EDTA·4Na	4	
	4-9	Inorganic salt medium (no-treated area)	—	100

Example 5 (evaluation of water culture of tomato seedlings)

[0053] Seeds of tomato "Momotaro" were sown in a box, and seedlings having 3 true leaves at the expansion period were used. For the respective compositions, the test was repeated 3 times. OKF2 (made by Otsuka Chemical Co., Ltd.) was diluted (as N:P:K = 260:149:247 (ppm)/OKF2 to be 538 times) as a NPK base and then resultant was added to a culturing solution. The present test was performed under conditions shown in Table 5. After 6 days from the start of the test, the culturing solution was sampled, and it was examined by means of RQ Flex (made by Merk) to obtain absorption efficiency of nitrate nitrogen. The each value as relative value shows an absorption amount of nitrate nitrogen at each treatment in the case of making NPK culturing solution to be control. Also, at 6 days after starting of test, SPAD value (SPAD 502 made by Minolta Co., Ltd.) representing green degree of a leaf were measured. When the control (Comparative product 5-8) made up to be 100, the relative value is compared. The results are shown in Table 5. Then, fertilizer composition of OKF2 (Otsuka Chemical Co., Ltd.) was as follows: N:P:K:Ca:Mg = 14:8:16:6:2.

Table 5

	Test No.	Blended raw materials	Concentration (ppm)	Result of the test	
				Efficiency of absorbing nitrogen	SPAD value
Invention product	5-1	Stearyl alcohol POE(20) sorbitan monooleate EDTA·4Na	50 150 20	207	109
	5-2	Stearyl alcohol POE(20) sorbitan monooleate	50 150	182	107
	5-3	Stearyl alcohol POE(20) sorbitan monooleate Lauryl sulfuric acid sodium salt EDTA·4Na	50 150 150 20	175	106
	5-4	Stearyl alcohol POE(20) sorbitan monooleate Lauryl sulfuric acid sodium salt	50 150 150	170	105
	5-5	Stearyl alcohol Lauryl sulfuric acid sodium salt EDTA·4Na	50 150 20	125	103
	5-6	Lauryl alcohol POE(20) sorbitan monooleate	50 150	134	104
	5-7	Cetyl alcohol POE(20) sorbitan monooleate	100 150	158	105
	5-8	Eicosanol POE(20) sorbitan monooleate	50 150	162	104
	5-9	Tetracosanol POE(20) sorbitan monooleate	50 150	146	103
	5-10	Lauryl alcohol POE(20) sorbitan monooleate Lauryl sulfuric acid sodium salt	50 150 150	128	102
	5-11	Cetyl alcohol POE(20) sorbitan monooleate Lauryl sulfuric acid sodium salt	50 150 150	149	104
	5-12	Eicosanol POE(20) sorbitan monooleate Lauryl sulfuric acid sodium salt	50 150 150	145	103
	5-13	Tetracosanol POE(20) sorbitan monooleate Lauryl sulfuric acid sodium salt	50 150 150	135	101
Comparative product	5-1	Ethanol POE(20) sorbitan monooleate	50 150	100	95
	5-2	Hexanol POE(20) sorbitan monooleate	50 150	97	97
	5-3	Decyl alcohol POE(20) sorbitan monooleate	50 150	95	96
	5-4	NPK culturing solution (control)	—	100	100

Example 6 (Test of soil-irrigating treatment for spinach: leaf vegetable)

[0054]

5 Species: "Esper"
Form of cultivation: a pot (diameter: 18 cm) for cultivation
Used soil: Kureha Engei Baido (horticultural soil made by Kureha Chemical Industry Co., Ltd.)
(Fertilizer components; N:P:K = 0.4:1.9:0.6 (g) per 1 kg of the soil)

10 [0055] Seeds of spinach "Esper" were directly sown on the soil. As the culturing soil, the horticultural soil (made by Kureha Chemical Industry Co., Ltd.) was used in an amount of 1.3 L (1.5 kg) per 1 pot (diameter of 18 cm). The number of repetition was made up to 18 pots. After 12 days from the sowing, treatment was started. At the intervals of 7 days, raw materials shown in Tables 6 and 7 were used to irrigate the soil 5 times in a treating amount of 150 ml/18 cm pot. After 6 days from the 5th irrigating treatment, weights of the plant and the SPAD value were examined. When the no-treated area (Comparative product 6-5) made up to be 100, the relative value is compared. The results are shown in
15 Tables 6 and 7. During the test period, fertilizer components such as nitrogen, phosphorous and potassium were not applied as additional fertilizer and the plant absorbed only nutrients contained in the soil. Then, composition ratio of NPK fertilizer as the additional fertilizer was follows: N:P:K:Mg = 17:9.5:18:3.5.

20

25

30

35

40

45

50

55

Table 6

Test No.	Blended raw materials	Concentration (ppm)	Result of the test	
			Weight	SPAD value
Invention product	6-1 Stearyl alcohol POE(20) sorbitan monooleic acid ester (RHEODOL TW-O120)	50 150	125	115
	6-2 Stearyl alcohol POE(6) sorbitan monooleic acid ester (RHEODOL TW-O106)	50 150	121	115
	6-3 Stearyl alcohol POE(20) sorbitan monostearic acid ester (RHEODOL TW-S120)	50 150	125	112
	6-4 Stearyl alcohol POE(20) sorbitan tristearic acid ester (RHEODOL TW-S320)	50 150	122	110
	6-5 Stearyl alcohol POE(20) sorbitan coconut oil fatty acid ester (RHEODOL TW-L120)	50 150	123	114
	6-6 Stearyl alcohol Sorbitan monostearic acid ester (RHEODOL SP-S10)	50 150	122	112
	6-7 Stearyl alcohol Sorbitan monooleic acid ester (RHEODOL SP-O10)	50 150	123	113
	6-8 Stearyl alcohol Sorbitan tristearic acid ester (RHEODOL SP-S30)	50 150	122	112
	6-9 Stearyl alcohol Sorbitan monopalmitic acid ester (RHEODOL SP-P10)	50 150	121	110
	6-10 Stearyl alcohol Sorbitan coconut oil fatty acid ester (RHEODOL SP-L10)	50 150	122	112
	6-11 Stearyl alcohol POE(30) sorbit tetraoleic acid ester (RHEODOL 430)	50 150	121	112
	6-12 Stearyl alcohol POE(40) sorbit tetraoleic acid ester (RHEODOL 440)	50 150	123	114
	6-13 Stearyl alcohol POE(60) sorbit tetraoleic acid ester (RHEODOL 460)	50 150	122	113
	6-14 Stearyl alcohol POE(140) monostearic acid polyethylene glycol (EMANON 3199)	50 150	122	114
	6-15 Stearyl alcohol POE(140) distearic acid polyethylene glycol (EMANON 3299)	50 150	121	111
	6-16 Stearyl alcohol POE(40) hardened castor oil (EMANON CH-40)	50 150	122	113
	6-17 Stearyl alcohol POE(80) hardened castor oil (EMANON CH-80)	50 150	121	113

- In the case of the testing, each of agents is used after forced-emulsifying by a home-mixer.
- Each of notes in parenthesis shows tradename made by Kao Corp.

Table 7

Test No.	Blended raw materials	Concentration (ppm)	Result of the test	
			Weight	SPAD value
6-18	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	120	112
	Oleic acid potassium salt (OS-SOAP)	150		
6-19	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	119	110
	Semi-hardened beef tallow fatty acid sodium salt (NS-SOAP)	150		
6-20	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	118	110
	Stearic acid sodium salt (SS-40N)	150		
6-21	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	117	111
	POE(4.5) lauryl ether acetic acid sodium salt (AKYPO RLM45NV)	150		
6-22	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	116	111
	POE(10) myristyl ether acetic acid sodium salt (AKYPO RLM100NV)	150		
6-23	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	115	112
	Lauryl phosphoric acid monoester sodium salt	150		
6-24	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	117	110
	Lauryl phosphoric acid diester sodium salt	150		
6-25	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	118	110
	POE(3) lauryl phosphoric acid monoester sodium salt	150		
6-26	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	117	111
	POE(4) lauryl ether (EMULGEN 104P)	75		
6-27	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	115	111
	POE(13) cetyl ether (EMULGEN 220)	75		
6-28	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	115	113
	POE(12) stearyl ether (EMULGEN 320P)	75		
6-29	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	116	112
	POE(9) octyl phenyl ether (EMULGEN 810)	75		
6-30	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	114	110
	POE(9) nonyl phenyl ether (EMULGEN 909)	75		
6-31	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	112	109
	POE(50) nonyl phenyl ether (EMULGEN 950)	75		
6-32	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	113	110
	Alkyl glucoside (C10/C12/C14) (MYDOL 12)	150		
6-33	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	112	107
	Alkyl glucoside (C9/C10/C11) (MYDOL 10)	150		
6-34	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	111	105
	Lauryl diethyl amine oxide (AMPHITOL 20N)	150		
6-35	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	111	110
	2-alkyl-N-carboxy methyl-N-hydroxy ethyl imidazolium betaine	150		
6-36	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	110	110
	Lauryl amido propyl betaine (AMPHITOL 20AB)	150		
6-37	C ₁₆ /C ₁₈ alcohol (C ₁₆ /C ₁₈ = 15/85)	50	109	107
	Lauryl dimethyl amino acetic acid betaine (AMPHITOL 24B)	150		
Comparative product	Decyl alcohol	50	75	69
	Lauryl sulfuric acid sodium salt (EMAL 0)	150		
	Hexacosanol	50	90	92
	Lauryl sulfuric acid sodium salt (EMAL 0)	150		
	Decyl alcohol	50	68	72
6-3	POE(9) nonyl phenyl ether (EMULGEN 909)	150		
	Hexacosanol	50	85	87
	POE(9) nonyl phenyl ether (EMULGEN 909)	150		
6-5	Water (no-treated area)	—	100	100

- In the case of the testing, each of agents is used after forced-emulsifying by a home-mixer.
- Each of notes in parenthesis shows tradename made by Kao Corp.

Example 7 (Test of soil-treating for rice-plant seedlings: treatment at the raising seedling period)

[0056] Unhulled rice seeds (species of Koshihikari) were sown into a box (60 × 30 cm) for raising seedling. At the period transferring from the hardening period to the greening period after germination, treatment was started in an treating amount of 500 ml/the box for raising seedling. Raw materials shown in Table 8 were used to irrigate the soil 3 times into soil. After the 3 treatments, that is, after 15 days from the sowing, the heights and the weights of the seedlings and the SPAD value were examined. When the no-treated area (Comparative product 7-6) made up to be 100, the relative value is compared. The results are shown in Table 8. As basal fertilizer, N component was applied in an amount of 0.5 g per the box for raising seedling. During the test period, fertilizer components were not applied as additional fertilizer and rice-plant seedlings absorbed only nutrients contained in the soil.

Table 8

Test No.	Blended raw materials	Concentration (ppm)	Result of the test		
			Height	Weight	SPAD value
Invention product	7-1 Stearyl alcohol POE(20) sorbitan monooleate	100 300	106	115	113
	7-2 Myristyl alcohol POE(20) sorbitan monooleate tartaric acid Na salt	100 300 50	107	107	108
	7-3 Tetracosanol POE(20) sorbitan monooleate	100 300	102	107	106
	7-4 Myristyl alcohol POE(20) sorbitan monooleate Oxamide	100 300 50	106	108	110
	7-5 Lauryl alcohol POE(20) sorbitan monolaurate	100 300	102	102	105
	7-6 Eicosanol POE(20) sorbitan monooleate	100 300	104	110	109
	7-7 Oleyl alcohol POE(20) sorbitan monooleate EDTA·4Na	100 300 50	108	119	116
	7-8 Stearyl alcohol Cetyl alcohol POE(20) sorbitan monooleate succinic acid	50 50 300 50	108	118	116
	7-9 Docosanol POE(20) sorbitan monooleate	100 300	103	108	108
	7-10 Myristyl alcohol POE(20) sorbitan monooleate	100 300	102	104	108
	7-11 Docosanol POE(20) sorbitan monooleate EDTA·4Na	100 300 50	106	116	110
	7-12 Cetyl alcohol POE(20) sorbitan monooleate	100 300	104	106	110
	7-13 Stearyl alcohol POE(20) sorbitan monooleate Malonic acid	100 300 50	109	119	115
	7-14 Cetyl alcohol POE(20) sorbitan monooleate EDTA·4Na	100 300 50	108	112	112
	7-15 Stearyl alcohol POE(20) sorbitan monooleate EDTA·4Na	100 300 50	110	121	118
	7-16 Oleyl alcohol POE(20) sorbitan monooleate	100 300	104	113	110
	7-17 Stearyl alcohol Cetyl alcohol POE(20) sorbitan monooleate	50 50 300	104	109	112
	7-18 Tetracosanol POE(20) sorbitan monooleate gluconic acid	100 300 50	104	110	109
	7-19 Eicosanol POE(20) sorbitan monooleate Ascorbic acid Na salt	100 300 50	108	118	114
	7-20 Lauryl alcohol POE(20) sorbitan monolaurate Fumaric acid	100 300 50	104	104	106
Comparative product	7-1 Ethanol POE(20) sorbitan monooleate	100 300	96	92	91
	7-2 Ethanol POE(20) sorbitan monooleate Heptonic acid Na salt	100 300 50	97	96	95
	7-3 Hexanol POE(20) sorbitan monooleate Malonic acid	100 300 50	93	96	89
	7-4 Hexanol POE(20) sorbitan monooleate	100 300	90	93	87
	7-5 Octyl alcohol POE(20) sorbitan monooleate	100 300	78	82	83
	7-6 Water (non-treated area)	—	100	100	100

Claims

1. Use of a mono-alcohol having 12 to 24 carbon atoms as a plant-activating agent.
- 5 2. Use as claimed in Claim 1, in which the plant-activating agent further comprises at least one compound selected from a surfactant, a fertilizer component and a chelating agent.
3. Use as claimed in Claim 2, in which the surfactant is selected from an ester group-containing nonionic surfactant, an ether group-containing nonionic surfactant having no nitrogen atom, an amphoteric surfactant, a carboxylic anionic surfactant and a phosphoric anionic surfactant.
- 10 4. A plant-activating composition comprising a mono-alcohol having 12 to 24 carbon atoms and one group selected from (i) a chelating agent, (ii) a surfactant and a chelating agent, (iii) a fertilizer component and a chelating agent and (iv) a surfactant, a fertilizer component and a chelating agent.
- 15 5. The composition as claimed in Claim 4, which comprises a mono-alcohol having 12 to 24 carbon atoms and at least one surfactant selected from an ester group-containing nonionic surfactant, an ether group-containing nonionic surfactant having no nitrogen atom, an amphoteric surfactant, a carboxylic anionic surfactant and a phosphoric anionic surfactant.
- 20 6. A plant-activating composition comprising a mono-alcohol having 12 to 24 carbon atoms and at least one surfactant selected from an ester group-containing nonionic surfactant, an ether group-containing nonionic surfactant having no nitrogen atom, an amphoteric surfactant, a carboxylic anionic surfactant and a phosphoric anionic surfactant.
- 25 7. The composition as claimed in Claim 6, which further comprises a fertilizer component.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 12 1907

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
P, X	WO 98 51148 A (WILEY ROBERT P ; MCKELVIE MARIA J (GB); FOLCHI BRUNO (IT); UNIROYAL) 19 November 1998 (1998-11-19) * page 1 - page 4 *	1-3	A01N31/02
X	DATABASE WPI Section Ch, Week 199806 Derwent Publications Ltd., London, GB; Class C03, AN 1998-052852 XP002130050 & CN 1 137 861 A (YU E), 18 December 1996 (1996-12-18) * abstract *	1,2,4	
Y		1-7	
X	US 4 169 716 A (ASHMEAD HARVEY H) 2 October 1979 (1979-10-02) * column 1, paragraphs 2,5,6 * * column 2, paragraph 3 * * column 3, paragraph 2 *	1 1-7	
X	US 5 385 750 A (ALEKSEJCZYK ROBERT A ET AL) 31 January 1995 (1995-01-31) * column 1, paragraph 3 * * column 2, paragraph 8 - paragraph 10 * * column 17, paragraph 6 * * claims 1,23 *	1-3,6,7 1-7	TECHNICAL FIELDS SEARCHED (Int.Cl.7) A01N
Y			
X	US 5 693 592 A (ILLINGWORTH JOHN) 2 December 1997 (1997-12-02) * column 1, paragraphs 1,6 * * column 2, line 6 * * column 2, line 48 - column 3, paragraph 11 *	1-3,6,7 1-7	
Y			
X	US 3 900 307 A (ABRAMITIS WALTER W) 19 August 1975 (1975-08-19) * column 1 * * column 2, line 32 - column 3, line 20 *	1-3,6,7 1-7	
Y			
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 9 February 2000	Examiner Bertrand, F
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons</p> <p>& : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.82 (P/C01)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 99 12 1907

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Y	GB 1 327 092 A (MURPHY CHEMICAL LTD) 15 August 1973 (1973-08-15) * page 1, line 11 - line 24 * * page 1, line 75 - page 2, line 82 * -----	2-7	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 9 February 2000	Examiner Bertrand, F
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03/82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 12 1907

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

09-02-2000

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9851148	A	19-11-1998	AU	7384398 A	08-12-1998
CN 1137861	A	18-12-1996	NONE		
US 4169716	A	02-10-1979	NONE		
US 5385750	A	31-01-1995	AU	4237793 A	13-12-1993
			CA	2135637 A	25-11-1993
			EP	0639943 A	01-03-1995
			JP	7508024 T	07-09-1995
			MX	9302843 A	28-02-1994
			WO	9322917 A	25-11-1993
US 5693592	A	02-12-1997	AU	704377 B	22-04-1999
			AU	5237096 A	05-12-1996
			NZ	286629 A	26-06-1998
US 3900307	A	19-08-1975	CA	1018791 A	11-10-1977
			JP	49126821 A	04-12-1974
GB 1327092	A	15-08-1973	AT	309480 B	15-07-1973
			BE	765774 A	15-10-1971
			CH	566944 A	30-09-1975
			CS	164886 B	28-11-1975
			DE	2119140 A	11-11-1971
			EG	10592 A	31-01-1976
			ES	389474 A	16-04-1974
			FR	2086482 A	31-12-1971
			IL	36475 A	29-11-1974
			IT	942314 B	20-03-1973
			NL	7105771 A	02-11-1971
			OA	3710 A	24-12-1971
			PH	9704 A	19-02-1976
			ZA	7101799 A	29-12-1971

EPO FORM P459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

